



## OCCURRENCE OF B<sub>1</sub> AFLATOXIN IN DIET AND M<sub>1</sub> AFLATOXIN IN BOVINE MILK

### OCORRÊNCIA DE AFLATOXINA B<sub>1</sub> NA DIETA E DE AFLATOXINA M<sub>1</sub> NO LEITE DE BOVINOS

ADRIANA FRIZZARIN<sup>1</sup>, THIAGO PEREIRA MOTTA<sup>1</sup>, THAMIRES MARTINS<sup>1</sup>, LIVIA CASTELANI<sup>1</sup>, HELOISA SOLDA DE AZEVEDO<sup>1</sup>,  
 CLÁUDIA RODRIGUES POZZI<sup>2</sup>

<sup>1</sup>Pós-Graduação em Produção Animal Sustentável, Instituto de Zootecnia (IZ), Agência Paulista de Tecnologia dos Agronegócios (APTA), Secretaria de Agricultura e Abastecimento do Estado de São Paulo (SAA), Rua Heitor Penteado, 56, Centro, CEP 13460-000, Nova Odessa, SP, Brasil. E-mail [adrianafrizzarin@hotmail.com](mailto:adrianafrizzarin@hotmail.com)

<sup>2</sup>Centro Avançado de Pesquisa Tecnológica do Agronegócio de Bovinos de Leite (CAPTA – Bovinos de Leite), IZ, APTA, SAA, Nova Odessa, SP, Brasil.

Ensuring food quality is one of the principles of food safety. Food for dairy cattle may be contaminated by fungi of the genus *Aspergillus*, which produce aflatoxins. The B<sub>1</sub> aflatoxin, when ingested by animals, is biotransformed in liver in several other toxic metabolites, including M<sub>1</sub> aflatoxin which is excreted in milk. M<sub>1</sub> aflatoxin has a carcinogenic effect, which the presence in milk poses a serious risk to public health because milk and dairy products are consumed mainly by children, pregnant women and elderly. The objective of this study was to detect the presence of B<sub>1</sub> aflatoxin in feed supplied to dairy cows and the presence of M<sub>1</sub> aflatoxin in milk. Samples were collected from complete diet (corn silage and concentrate) from a batch of 15 lactating cows from a dairy farm in the Campinas region. Two samples of diets were collected directly into the troughs in intervals of 24 hours at every 15 days, totalizing a period of 45 days. Milk samples of those cows were collected 24 hours after diet collection, directly from sample valves in the glass jars. B<sub>1</sub> and M<sub>1</sub> aflatoxins were detected by the technique of High Performance Liquid Chromatography after extraction and purification on immunoaffinity columns. From the 40 samples of diets evaluated, 40% were contaminated with B<sub>1</sub> aflatoxin, and the levels found ranged from 1.93 to 43.78 µg/Kg. One sample showed result higher than the maximum recommended for grain and animal feed in Brazil (20 µg/Kg). From the 75 milk samples analyzed, the presence of M<sub>1</sub> aflatoxin was detected in 13.3% with levels ranging from 0.03 to 0.16 µg/L, not exceeding the maximum permitted for marketing in the country of 0.5 µg/L, however 80% of contaminated samples had values above the maximum permissible levels of 0.05 µg/L, value found among countries with abundant milk production... The presence of aflatoxins highlights the importance of monitoring the production, the storage and the importance of handling food and ingredients intended for dairy cattle to prevent the presence of B<sub>1</sub> aflatoxin and consequently, M<sub>1</sub> aflatoxin in milk.

Key words: B<sub>1</sub> aflatoxin; M<sub>1</sub> aflatoxin; diet, lactating cows.